



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by L. C. WALKER, A.M., Professor of Mathematics, Petaluma High School, Petaluma, Cal.

Equation of the cordiod is $r=a(1+\cos\theta)$, and its complete perimeter is

$$S=2\int_0^\pi [a^2(1+\cos\theta)^2+a^2\sin^2\theta]d\theta=4a\int_0^\pi \cos\frac{1}{2}\theta d\theta=8a,$$

the arc S being measured from the point where the curve crosses the initial line.

Let θ' be the point of the first division; then

$$2a\int_0^{\theta'} \cos\frac{1}{2}\theta d\theta=4a\sin\frac{1}{2}\theta'=a;$$

from which $\theta_1, \theta_7=\pm 2\sin^{-1}\frac{1}{4}$. Similarly, $\theta_2, \theta_6=\pm 2\sin^{-1}\frac{1}{2}$, $\theta_3, \theta_5=\pm 2\sin^{-1}\frac{3}{4}$, $\theta_4=\pi$, and $\theta_8=0$.

Excellent solutions were received from G. B. M. ZERR, and F. P. MATZ.

109. Proposed by J. SCHEFFER, A. M., Hagerstown. Md.

Find the latitude of the place where the sun's center remains above the horizon for a hundred successive days.

Solution by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy. Defiance College, Defiance, O.

I. The solar phenomenon in question will begin May 2 and end August 10. For May 2 at Washington mean noon, according to the Director of United States Nautical Almanac Office, the declination of the sun is $\delta=+15^\circ 18' 54.7''$. Representing the hour-angle of the sun by h , we have $\cosh=-\tan\delta\tan\theta\dots(1)$.

Since there is to be no setting of the sun, we may put $\cosh=-1$.

$\therefore \lambda=\tan^{-1}\left(\frac{1}{\tan\delta}\right)=\tan^{-1}(\cot\delta)=90^\circ-\delta=74^\circ 41' 5.3''$, which is the terrestrial latitude of the place required. This solar phenomenon may be observed on Melville Island.

II. The celestial longitude of the sun for mean noon Washington can be determined in various ways to be about $\psi=41^\circ 34' 25''$; the obliquity of the ecliptic may be taken $\omega=23^\circ 27' 8''$; then $\lambda=90-\delta=90-\sin^{-1}(\sin\psi\times\sin\omega),=74^\circ 43' 16''$.

Also solved by G. B. M. ZERR, and H. C. WHITAKER.

110. Proposed by E. W. MORELL, South Trowbridge, Vt.

If a and b be the sides of a triangle, A and B the angles opposite, then will $\log b - \log a = \cos 3A - \cos 2B + \frac{1}{2}(\cos 4A - \cos 4B) + \frac{1}{8}(\cos 6A - \cos 6B) + \dots$

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

$$1 + \cos 2A + \frac{1}{2}\cos 4A + \frac{1}{8}\cos 6A + \dots = -\frac{1}{2}\log(2 - 2\cos 2A) = -\log(2\sin A).$$

$$1 + \cos 2B + \frac{1}{2}\cos 4B + \frac{1}{8}\cos 6B + \dots = -\log(2\sin B).$$

See Trigonometry for the summation of these series.

$$\therefore \cos 2A - \cos 2B + \frac{1}{2}(\cos 4A - \cos 4B) + \dots = \log(\sin B/\sin A) = \log(b/a).$$

Also solved by F. P. MATZ, and J. SCHEFFER.